


May 11, 2014

## **NEW PRODUCTS**

## Native & Mass-Labelled TBBA and TBPAn

Although structurally similar, 2,3,4,5-tetrabromobenzoic acid (TBBA) and tetrabromophthalic anhydride (TBPAn) have very different origins in environmental samples. TBBA does not have a known commercial application; it appears to be arising as a metabolite of 2-ethylhexyl-2,3,4,5-tetrabromobenzoate (a component of several popular flame retardant mixtures). Conversely, TBPAn has been marketed as both an additive and reactive flame retardant. TBPAn appears to be most commonly used as a reactive intermediate during the production of saturated and unsaturated polyesters, polyols, esters, and imides. In order to aid researchers in the accurate quantification of these compounds in environmental samples, Wellington has synthesized the following native and mass-labelled reference standards:



Catalogue Number	Product (methanol)	Qty	Conc
ТВВА	2,3,4,5-Tetrabromobenzoic acid	1.2 ml	50 μg/ml
МТВВА	2,3,4,5-Tetrabromobenzoic acid [13C <sub>6</sub> ring]	1.2 ml	50 μg/ml

Catalogue Number	Product (toluene)	Qty	Conc
MeTBBA	Methyl-2,3,4,5-tetrabromobenzoate	1.2 ml	50 μg/ml
TBPAn	Tetrabromophthalic anhydride	1.2 ml	50 μg/ml
MTBPAn	Tetrabromo[13C <sub>6</sub> ]phthalic anhydride	1.2 ml	50 μg/ml



Please contact your local distributor or info@well-labs.com for pricing and delivery.

Visit our website (www.well-labs.com) for a complete listing of our new products.